























#	Template	Alignment Coverage	3D Model	Confidence	% i.d.	Template Information
1	<a href="#">c2e85B_</a>	 Alignment		100.0	99	<b>PDB header:</b> hydrolase <b>Chain:</b> B: <b>PDB Molecule:</b> hydrogenase 3 maturation protease; <b>PDBTitle:</b> crystal structure of the hydrogenase 3 maturation protease
2	<a href="#">d1cfza_</a>	 Alignment		100.0	18	<b>Fold:</b> Phosphorylase/hydrolase-like <b>Superfamily:</b> HybD-like <b>Family:</b> Hydrogenase maturing endopeptidase HybD
3	<a href="#">c3pu6A_</a>	 Alignment		100.0	33	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> uncharacterized protein; <b>PDBTitle:</b> the crystal structure of an uncharacterized protein from wolinetla2 succinogenes
4	<a href="#">d1c8ba_</a>	 Alignment		96.5	12	<b>Fold:</b> Phosphorylase/hydrolase-like <b>Superfamily:</b> HybD-like <b>Family:</b> Germination protease
5	<a href="#">c3neaA_</a>	 Alignment		94.4	19	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> peptidyl-trna hydrolase; <b>PDBTitle:</b> crystal structure of peptidyl-trna hydrolase from francisella2 tularensis
6	<a href="#">d2ptha_</a>	 Alignment		91.6	19	<b>Fold:</b> Phosphorylase/hydrolase-like <b>Superfamily:</b> Peptidyl-tRNA hydrolase-like <b>Family:</b> Peptidyl-tRNA hydrolase-like
7	<a href="#">d1lssa_</a>	 Alignment		82.2	19	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> Potassium channel NAD-binding domain
8	<a href="#">c3l4bG_</a>	 Alignment		81.9	11	<b>PDB header:</b> transport protein <b>Chain:</b> G: <b>PDB Molecule:</b> trka k+ channel protien tm1088b; <b>PDBTitle:</b> crystal structure of an octomeric two-subunit trka k+ channel ring2 gating assembly, tm1088a:tm1088b, from thermotoga maritima
9	<a href="#">c2z2iA_</a>	 Alignment		81.7	16	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> peptidyl-trna hydrolase; <b>PDBTitle:</b> crystal structure of peptidyl-trna hydrolase from mycobacterium2 tuberculosis
10	<a href="#">d2vapa1</a>	 Alignment		78.4	13	<b>Fold:</b> Tubulin nucleotide-binding domain-like <b>Superfamily:</b> Tubulin nucleotide-binding domain-like <b>Family:</b> Tubulin, GTPase domain
11	<a href="#">c3eywA_</a>	 Alignment		76.7	23	<b>PDB header:</b> transport protein <b>Chain:</b> A: <b>PDB Molecule:</b> c-terminal domain of glutathione-regulated potassium-efflux <b>PDBTitle:</b> crystal structure of the c-terminal domain of e. coli kefc in complex2 with keff

12	<a href="#">c1w5fA_</a>	Alignment		76.5	15	<b>PDB header:</b> cell division <b>Chain:</b> A: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> ftsz, t7 mutated, domain swapped (t. maritima)
13	<a href="#">d1w5fa1</a>	Alignment		76.1	15	<b>Fold:</b> Tubulin nucleotide-binding domain-like <b>Superfamily:</b> Tubulin nucleotide-binding domain-like <b>Family:</b> Tubulin, GTPase domain
14	<a href="#">c1w59B_</a>	Alignment		73.0	13	<b>PDB header:</b> cell division <b>Chain:</b> B: <b>PDB Molecule:</b> cell division protein ftsz homolog 1; <b>PDBTitle:</b> ftsz dimer, empty (m. jannaschii)
15	<a href="#">c3rh0A_</a>	Alignment		71.5	27	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> arsenate reductase; <b>PDBTitle:</b> corynebacterium glutamicum mycothiol/mycoredoxin1-dependent arsenate2 reductase cg_arsc2
16	<a href="#">c3vh3A_</a>	Alignment		71.1	24	<b>PDB header:</b> metal binding protein/protein transport <b>Chain:</b> A: <b>PDB Molecule:</b> ubiquitin-like modifier-activating enzyme atg7; <b>PDBTitle:</b> crystal structure of atg7ctd-atg8 complex
17	<a href="#">d2g2ca1</a>	Alignment		66.3	13	<b>Fold:</b> Molybdenum cofactor biosynthesis proteins <b>Superfamily:</b> Molybdenum cofactor biosynthesis proteins <b>Family:</b> MogA-like
18	<a href="#">c2pjka_</a>	Alignment		66.2	8	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> 178aa long hypothetical molybdenum cofactor <b>PDBTitle:</b> structure of hypothetical molybdenum cofactor biosynthesis2 protein b from sulfobobus tokodaii
19	<a href="#">c3ic5A_</a>	Alignment		62.2	19	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> putative saccharopine dehydrogenase; <b>PDBTitle:</b> n-terminal domain of putative saccharopine dehydrogenase from ruegeria2 pomeroyi.
20	<a href="#">c3gucB_</a>	Alignment		60.4	23	<b>PDB header:</b> transferase <b>Chain:</b> B: <b>PDB Molecule:</b> ubiquitin-like modifier-activating enzyme 5; <b>PDBTitle:</b> human ubiquitin-activating enzyme 5 in complex with amppnp
21	<a href="#">c2zklA_</a>	Alignment	not modelled	60.4	14	<b>PDB header:</b> isomerase <b>Chain:</b> A: <b>PDB Molecule:</b> capsular polysaccharide synthesis enzyme cap5f; <b>PDBTitle:</b> crystal structure of capsular polysaccharide assembling protein capf2 from staphylococcus aureus
22	<a href="#">c3jviA_</a>	Alignment	not modelled	56.9	12	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> protein tyrosine phosphatase; <b>PDBTitle:</b> product state mimic crystal structure of protein tyrosine phosphatase2 from entamoeba histolytica
23	<a href="#">d1ryba_</a>	Alignment	not modelled	56.0	21	<b>Fold:</b> Phosphorylase/hydrolase-like <b>Superfamily:</b> Peptidyl-tRNA hydrolase-like <b>Family:</b> Peptidyl-tRNA hydrolase-like
24	<a href="#">c2yxba_</a>	Alignment	not modelled	55.2	11	<b>PDB header:</b> isomerase <b>Chain:</b> A: <b>PDB Molecule:</b> coenzyme b12-dependent mutase; <b>PDBTitle:</b> crystal structure of the methylmalonyl-coa mutase alpha-subunit from2 aeropyrum pernix
25	<a href="#">d2f1ka2</a>	Alignment	not modelled	51.9	17	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
26	<a href="#">c3ckyA_</a>	Alignment	not modelled	50.2	13	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> 2-hydroxymethyl glutarate dehydrogenase; <b>PDBTitle:</b> structural and kinetic properties of a beta-hydroxyacid dehydrogenase2 involved in nicotinate fermentation
27	<a href="#">c3rfqC_</a>	Alignment	not modelled	49.8	17	<b>PDB header:</b> biosynthetic protein <b>Chain:</b> C: <b>PDB Molecule:</b> pterin-4-alpha-carbinolamine dehydratase moab2; <b>PDBTitle:</b> crystal structure of pterin-4-alpha-carbinolamine dehydratase moab22 from mycobacterium marinum
						<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> H: <b>PDB Molecule:</b> similar to 2-dehydropantoate 2-

28	<a href="#">c3g17H_</a>	Alignment	not modelled	49.6	13	reductase; <b>PDBTitle:</b> structure of putative 2-dehydropantoate 2-reductase from2 staphylococcus aureus <b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> crystal structure of mycobacterium tuberculosis low2 molecular protein tyrosine phosphatase (mptpa) at 1.9a3 resolution <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
29	<a href="#">c1u2pA_</a>	Alignment	not modelled	49.2	12	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> crystal structure of mycobacterium tuberculosis low2 molecular protein tyrosine phosphatase (mptpa) at 1.9a3 resolution <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
30	<a href="#">d1mv8a2</a>	Alignment	not modelled	49.1	21	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> crystal structure of mycobacterium tuberculosis low2 molecular protein tyrosine phosphatase (mptpa) at 1.9a3 resolution <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
31	<a href="#">c3gznB_</a>	Alignment	not modelled	49.0	26	<b>PDB header:</b> protein binding/ligase <b>Chain:</b> B: <b>PDB Molecule:</b> nedd8-activating enzyme e1 catalytic subunit; <b>PDBTitle:</b> structure of nedd8-activating enzyme in complex with nedd82 and mln4924 <b>Fold:</b> Phosphotyrosine protein phosphatases I-like <b>Superfamily:</b> Phosphotyrosine protein phosphatases I <b>Family:</b> Low-molecular-weight phosphotyrosine protein phosphatases
32	<a href="#">d1y1la_</a>	Alignment	not modelled	48.7	19	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> crystal structure of the s. aureus protein tyrosine phosphatase ptpa <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
33	<a href="#">c3rofa_</a>	Alignment	not modelled	48.5	12	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> crystal structure of the s. aureus protein tyrosine phosphatase ptpa <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
34	<a href="#">c2z2vA_</a>	Alignment	not modelled	47.2	26	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> hypothetical protein ph1688; <b>PDBTitle:</b> crystal structure of l-lysine dehydrogenase from2 hyperthermophilic archaeon pyrococcus horikoshii <b>Fold:</b> Molybdenum cofactor biosynthesis proteins <b>Superfamily:</b> Molybdenum cofactor biosynthesis proteins <b>Family:</b> MogA-like
35	<a href="#">d1mkza_</a>	Alignment	not modelled	46.7	16	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> ketopantoate reductase; <b>PDBTitle:</b> crystal structure of escherichia coli ketopantoate2 reductase in a ternary complex with nadp+ and pantoate <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
36	<a href="#">d1txga2</a>	Alignment	not modelled	45.4	17	<b>PDB header:</b> cell cycle <b>Chain:</b> A: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> the structure of ftsz from bacillus subtilis at 1.7a2 resolution <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
37	<a href="#">c2vxyA_</a>	Alignment	not modelled	44.8	16	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> ketopantoate reductase; <b>PDBTitle:</b> crystal structure of escherichia coli ketopantoate2 reductase in a ternary complex with nadp+ and pantoate <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
38	<a href="#">c2ofpB_</a>	Alignment	not modelled	44.8	17	<b>PDB header:</b> oxidoreductase <b>Chain:</b> D: <b>PDB Molecule:</b> prephenate dehydrogenase; <b>PDBTitle:</b> crystal structure of synechocystis arogonate dehydrogenase <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
39	<a href="#">c2f1kD_</a>	Alignment	not modelled	44.1	17	<b>PDB header:</b> oxidoreductase <b>Chain:</b> D: <b>PDB Molecule:</b> prephenate dehydrogenase; <b>PDBTitle:</b> crystal structure of synechocystis arogonate dehydrogenase <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
40	<a href="#">d2ngra3</a>	Alignment	not modelled	44.0	15	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> putative oxidoreductase; <b>PDBTitle:</b> crystal structure of a putative oxidoreductase from mycobacterium2 avium 104 <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
41	<a href="#">d1q0qa2</a>	Alignment	not modelled	43.8	20	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> structure of a protein tyrosine phosphatase <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
42	<a href="#">d1ks9a2</a>	Alignment	not modelled	43.5	17	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> structure of a protein tyrosine phosphatase <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
43	<a href="#">c3qhaB_</a>	Alignment	not modelled	43.4	33	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight protein-tyrosine- <b>PDBTitle:</b> structure of a protein tyrosine phosphatase <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
44	<a href="#">c2fekA_</a>	Alignment	not modelled	43.2	20	<b>PDB header:</b> transport protein <b>Chain:</b> A: <b>PDB Molecule:</b> putative glutathione-regulated potassium-efflux system <b>PDBTitle:</b> crystal structure of trka domain of putative glutathione-regulated2 potassium-efflux kefb from vibrio parahaemolyticus <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
45	<a href="#">c3c85A_</a>	Alignment	not modelled	42.9	21	<b>PDB header:</b> structural protein <b>Chain:</b> A: <b>PDB Molecule:</b> molybdopterin biosynthesis enzyme, moab; <b>PDBTitle:</b> crystal structure of the molybdopterin biosynthesis enzyme moab2 (ttha0341) from thermus thermophilus hb8 <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
46	<a href="#">c2is8A_</a>	Alignment	not modelled	42.4	15	<b>PDB header:</b> cell cycle <b>Chain:</b> 1: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> aquifex aeolicus ftsz <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
47	<a href="#">c2r6r1_</a>	Alignment	not modelled	42.2	13	<b>PDB header:</b> cell cycle <b>Chain:</b> 1: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> aquifex aeolicus ftsz <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
48	<a href="#">d1pgja2</a>	Alignment	not modelled	40.6	23	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight phosphotyrosine protein phosphatase; <b>PDBTitle:</b> crystal structure of tt1001 protein from thermus thermophilus hb8 <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
49	<a href="#">c2cwaA_</a>	Alignment	not modelled	40.6	14	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> low molecular weight phosphotyrosine protein phosphatase; <b>PDBTitle:</b> crystal structure of tt1001 protein from thermus thermophilus hb8 <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
50	<a href="#">d1t2da1</a>	Alignment	not modelled	40.3	24	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> 2-dehydropantoate 2-reductase; <b>PDBTitle:</b> ketopantoate reductase from escherichia coli <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
51	<a href="#">c1ks9A_</a>	Alignment	not modelled	40.0	17	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> 2-dehydropantoate 2-reductase; <b>PDBTitle:</b> ketopantoate reductase from escherichia coli <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
52	<a href="#">d1j5pa4</a>	Alignment	not modelled	39.3	40	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> 2-dehydropantoate 2-reductase; <b>PDBTitle:</b> ketopantoate reductase from escherichia coli <b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
						<b>Fold:</b> NAD(P)-binding Rossmann-fold domains

53	<a href="#">d1n1ea2</a>	Alignment	not modelled	38.9	20	<b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
54	<a href="#">c2vawA</a>	Alignment	not modelled	38.3	18	<b>PDB header:</b> cell cycle <b>Chain:</b> A: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> ftsz pseudomonas aeruginosa gdp
55	<a href="#">d1jlja</a>	Alignment	not modelled	38.0	14	<b>Fold:</b> Molybdenum cofactor biosynthesis proteins <b>Superfamily:</b> Molybdenum cofactor biosynthesis proteins <b>Family:</b> MogA-like
56	<a href="#">d1vl0a</a>	Alignment	not modelled	37.9	9	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> Tyrosine-dependent oxidoreductases
57	<a href="#">c3vh1A</a>	Alignment	not modelled	37.2	13	<b>PDB header:</b> metal binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> ubiquitin-like modifier-activating enzyme atg7; <b>PDBTitle:</b> crystal structure of saccharomyces cerevisiae atg7 (1-595)
58	<a href="#">c3lcmB</a>	Alignment	not modelled	36.9	9	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> putative oxidoreductase; <b>PDBTitle:</b> crystal structure of smu.1420 from streptococcus mutans ua159
59	<a href="#">d1yovb1</a>	Alignment	not modelled	36.9	26	<b>Fold:</b> Activating enzymes of the ubiquitin-like proteins <b>Superfamily:</b> Activating enzymes of the ubiquitin-like proteins <b>Family:</b> Ubiquitin activating enzymes (UBA)
60	<a href="#">c3dojA</a>	Alignment	not modelled	36.3	17	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> dehydrogenase-like protein; <b>PDBTitle:</b> structure of glyoxylate reductase 1 from arabidopsis2 (atglyr1)
61	<a href="#">c1nhqA</a>	Alignment	not modelled	35.9	16	<b>PDB header:</b> oxidoreductase (h2o2(a)) <b>Chain:</b> A: <b>PDB Molecule:</b> nadh peroxidase; <b>PDBTitle:</b> crystallographic analyses of nadh peroxidase cys42ala and cys42ser2 mutants: active site structure, mechanistic implications, and an3 unusual environment of arg303
62	<a href="#">c2gf2B</a>	Alignment	not modelled	35.6	27	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> 3-hydroxyisobutyrate dehydrogenase; <b>PDBTitle:</b> crystal structure of human hydroxyisobutyrate dehydrogenase
63	<a href="#">c2rhoB</a>	Alignment	not modelled	34.4	16	<b>PDB header:</b> cell cycle <b>Chain:</b> B: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> synthetic gene encoded bacillus subtilis ftsz ncs dimer with2 bound gdp and gtp-gamma-s
64	<a href="#">c1a5za</a>	Alignment	not modelled	33.8	23	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> l-lactate dehydrogenase; <b>PDBTitle:</b> lactate dehydrogenase from thermotoga maritima (tml dh)
65	<a href="#">c1vkzA</a>	Alignment	not modelled	33.4	22	<b>PDB header:</b> ligase <b>Chain:</b> A: <b>PDB Molecule:</b> phosphoribosylamine--glycine ligase; <b>PDBTitle:</b> crystal structure of phosphoribosylamine--glycine ligase (tml250) from2 thermotoga maritima at 2.30 a resolution
66	<a href="#">c1ojua</a>	Alignment	not modelled	33.2	19	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> malate dehydrogenase; <b>PDBTitle:</b> 2.8 a resolution structure of malate dehydrogenase from2 archaeoglobus fulgidus in complex with etheno-nad.
67	<a href="#">d5pnta</a>	Alignment	not modelled	33.1	14	<b>Fold:</b> Phosphotyrosine protein phosphatases I-like <b>Superfamily:</b> Phosphotyrosine protein phosphatases I <b>Family:</b> Low-molecular-weight phosphotyrosine protein phosphatases
68	<a href="#">c2cduB</a>	Alignment	not modelled	33.0	13	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> nadh oxidase; <b>PDBTitle:</b> the crystal structure of water-forming nad(p)h oxidase from2 lactobacillus sanfranciscensis
69	<a href="#">d1a5za1</a>	Alignment	not modelled	32.5	23	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
70	<a href="#">c2uyyD</a>	Alignment	not modelled	31.8	13	<b>PDB header:</b> cytokine <b>Chain:</b> D: <b>PDB Molecule:</b> n-pac protein; <b>PDBTitle:</b> structure of the cytokine-like nuclear factor n-pac
71	<a href="#">c2h1fB</a>	Alignment	not modelled	31.6	16	<b>PDB header:</b> transferase <b>Chain:</b> B: <b>PDB Molecule:</b> lipopolysaccharide heptosyltransferase-1; <b>PDBTitle:</b> e. coli heptosyltransferase waac with adp
72	<a href="#">c2gi4A</a>	Alignment	not modelled	31.5	12	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> possible phosphotyrosine protein phosphatase; <b>PDBTitle:</b> solution structure of the low molecular weight protein2 tyrosine phosphatase from campylobacter jejuni.
73	<a href="#">d1nhpa1</a>	Alignment	not modelled	30.9	19	<b>Fold:</b> FAD/NAD(P)-binding domain <b>Superfamily:</b> FAD/NAD(P)-binding domain <b>Family:</b> FAD/NAD-linked reductases, N-terminal and central domains
74	<a href="#">d1ofua1</a>	Alignment	not modelled	30.4	18	<b>Fold:</b> Tubulin nucleotide-binding domain-like <b>Superfamily:</b> Tubulin nucleotide-binding domain-like <b>Family:</b> Tubulin, GTPase domain
75	<a href="#">c1zfnA</a>	Alignment	not modelled	29.9	14	<b>PDB header:</b> transferase <b>Chain:</b> A: <b>PDB Molecule:</b> adenyllyltransferase thif; <b>PDBTitle:</b> structural analysis of escherichia coli thif
76	<a href="#">c3fvwA</a>	Alignment	not modelled	29.7	15	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> putative nad(p)h-dependent fmh reductase; <b>PDBTitle:</b> crystal structure of the q8dwd8_strmu protein from2 streptococcus mutans. northeast structural genomics3 consortium target smr99.
77	<a href="#">c1zggA</a>	Alignment	not modelled	29.7	31	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> putative low molecular weight protein-tyrosine- <b>PDBTitle:</b> solution structure of a low molecular weight protein2 tyrosine phosphatase from bacillus subtilis

78	<a href="#">dljw9b_</a>	Alignment	not modelled	28.5	11	<b>Fold:</b> Activating enzymes of the ubiquitin-like proteins <b>Superfamily:</b> Activating enzymes of the ubiquitin-like proteins <b>Family:</b> Molybdenum cofactor biosynthesis protein MoeB
79	<a href="#">dlc1da1</a>	Alignment	not modelled	28.4	28	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> Aminoacid dehydrogenase-like, C-terminal domain
80	<a href="#">c3ghyA_</a>	Alignment	not modelled	28.2	20	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> ketopantoate reductase protein; <b>PDBTitle:</b> crystal structure of a putative ketopantoate reductase from ralstonia2 solanacearum molk2
81	<a href="#">dlvzfa1</a>	Alignment	not modelled	28.0	22	<b>Fold:</b> Flavodoxin-like <b>Superfamily:</b> SGNH hydrolase <b>Family:</b> TAP-like
82	<a href="#">dlguza1</a>	Alignment	not modelled	27.6	26	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
83	<a href="#">dlrq2a1</a>	Alignment	not modelled	27.5	16	<b>Fold:</b> Tubulin nucleotide-binding domain-like <b>Superfamily:</b> Tubulin nucleotide-binding domain-like <b>Family:</b> Tubulin, GTPase domain
84	<a href="#">c3oc4A_</a>	Alignment	not modelled	27.3	19	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> oxidoreductase, pyridine nucleotide-disulfide family; <b>PDBTitle:</b> crystal structure of a pyridine nucleotide-disulfide family2 oxidoreductase from the enterococcus faecalis v583
85	<a href="#">c3qjaA_</a>	Alignment	not modelled	26.6	12	<b>PDB header:</b> lyase <b>Chain:</b> A: <b>PDB Molecule:</b> indole-3-glycerol phosphate synthase; <b>PDBTitle:</b> crystal structure of the mycobacterium tuberculosis indole-3-glycerol2 phosphate synthase (trpc) in apo form
86	<a href="#">dljf8a_</a>	Alignment	not modelled	26.5	19	<b>Fold:</b> Phosphotyrosine protein phosphatases I-like <b>Superfamily:</b> Phosphotyrosine protein phosphatases I <b>Family:</b> Low-molecular-weight phosphotyrosine protein phosphatases
87	<a href="#">clofuB_</a>	Alignment	not modelled	26.5	18	<b>PDB header:</b> bacterial cell division inhibitor <b>Chain:</b> B: <b>PDB Molecule:</b> cell division protein ftsz; <b>PDBTitle:</b> crystal structure of sul:fts from pseudomonas aeruginosa
88	<a href="#">dld1qa_</a>	Alignment	not modelled	25.9	15	<b>Fold:</b> Phosphotyrosine protein phosphatases I-like <b>Superfamily:</b> Phosphotyrosine protein phosphatases I <b>Family:</b> Low-molecular-weight phosphotyrosine protein phosphatases
89	<a href="#">d2bli1a1</a>	Alignment	not modelled	25.4	22	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> Tyrosine-dependent oxidoreductases
90	<a href="#">c2nvuB_</a>	Alignment	not modelled	25.3	27	<b>PDB header:</b> protein turnover, ligase <b>Chain:</b> B: <b>PDB Molecule:</b> maltose binding protein/nedd8-activating enzyme <b>PDBTitle:</b> structure of appbp1-uba3~nedd8-nedd8-mgatp-ubc12(c111a), a2 trapped ubiquitin-like protein activation complex
91	<a href="#">dluz5a3</a>	Alignment	not modelled	25.1	15	<b>Fold:</b> Molybdenum cofactor biosynthesis proteins <b>Superfamily:</b> Molybdenum cofactor biosynthesis proteins <b>Family:</b> MoeA central domain-like
92	<a href="#">c3hp7A_</a>	Alignment	not modelled	24.8	15	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> hemolysin, putative; <b>PDBTitle:</b> putative hemolysin from streptococcus thermophilus.
93	<a href="#">d2bfbdb2</a>	Alignment	not modelled	24.7	21	<b>Fold:</b> TK C-terminal domain-like <b>Superfamily:</b> TK C-terminal domain-like <b>Family:</b> Branched-chain alpha-keto acid dehydrogenase beta-subunit, C-terminal-domain
94	<a href="#">dlouja1</a>	Alignment	not modelled	24.6	19	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> LDH N-terminal domain-like
95	<a href="#">c3lp8A_</a>	Alignment	not modelled	24.0	19	<b>PDB header:</b> ligase <b>Chain:</b> A: <b>PDB Molecule:</b> phosphoribosylamine-glycine ligase; <b>PDBTitle:</b> crystal structure of phosphoribosylamine-glycine ligase from2 ehrlichia chaffeensis
96	<a href="#">c3l6dB_</a>	Alignment	not modelled	23.9	16	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> putative oxidoreductase; <b>PDBTitle:</b> crystal structure of putative oxidoreductase from pseudomonas putida2 kt2440
97	<a href="#">c3grzA_</a>	Alignment	not modelled	23.9	21	<b>PDB header:</b> transferase <b>Chain:</b> A: <b>PDB Molecule:</b> ribosomal protein l11 methyltransferase; <b>PDBTitle:</b> crystal structure of ribosomal protein l11 methylase from2 lactobacillus delbrueckii subsp. bulgaricus
98	<a href="#">dldg9a_</a>	Alignment	not modelled	23.8	18	<b>Fold:</b> Phosphotyrosine protein phosphatases I-like <b>Superfamily:</b> Phosphotyrosine protein phosphatases I <b>Family:</b> Low-molecular-weight phosphotyrosine protein phosphatases
99	<a href="#">c2dc1A_</a>	Alignment	not modelled	22.8	21	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> l-aspartate dehydrogenase; <b>PDBTitle:</b> crystal structure of l-aspartate dehydrogenase from2 hyperthermophilic archaeon archaeoglobus fulgidus
100	<a href="#">dl00sa2</a>	Alignment	not modelled	22.7	21	<b>Fold:</b> Aminoacid dehydrogenase-like, N-terminal domain <b>Superfamily:</b> Aminoacid dehydrogenase-like, N-terminal domain <b>Family:</b> Malic enzyme N-domain
101	<a href="#">dly0ya2</a>	Alignment	not modelled	22.6	18	<b>Fold:</b> Phosphorylase/hydrolase-like <b>Superfamily:</b> Zn-dependent exopeptidases <b>Family:</b> Bacterial dinuclear zinc exopeptidases
102	<a href="#">c2wmyH_</a>	Alignment	not modelled	22.5	14	<b>PDB header:</b> hydrolase <b>Chain:</b> H: <b>PDB Molecule:</b> putative acid phosphatase wzb; <b>PDBTitle:</b> crystal structure of the tyrosine phosphatase wzb from2 escherichia coli k30 in complex with sulphate.
103	<a href="#">c2g1uA_</a>	Alignment	not modelled	22.3	15	<b>PDB header:</b> transport protein <b>Chain:</b> A: <b>PDB Molecule:</b> hypothetical protein tm1088a; <b>PDBTitle:</b> crystal structure of a putative transport protein (tm1088a) from2 thermotoga maritima at 1.50 a resolution
						<b>PDB header:</b> oxidoreductase



104	<a href="#">c3prjB_</a>	Alignment	not modelled	21.6	16	<b>Chain:</b> B: <b>PDB Molecule:</b> udp-glucose 6-dehydrogenase; <b>PDBTitle:</b> role of packing defects in the evolution of allostery and induced fit2 in human udp-glucose dehydrogenase.
105	<a href="#">d1a53a_</a>	Alignment	not modelled	21.5	18	<b>Fold:</b> TIM beta/alpha-barrel <b>Superfamily:</b> Ribulose-phosphate binding barrel <b>Family:</b> Tryptophan biosynthesis enzymes
106	<a href="#">d1i36a2</a>	Alignment	not modelled	21.3	12	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
107	<a href="#">d1v8da_</a>	Alignment	not modelled	21.2	14	<b>Fold:</b> TTHA0583/YokD-like <b>Superfamily:</b> TTHA0583/YokD-like <b>Family:</b> Hypothetical protein TT1679
108	<a href="#">d1leha1</a>	Alignment	not modelled	21.1	13	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> Aminoacid dehydrogenase-like, C-terminal domain
109	<a href="#">d1a4ia1</a>	Alignment	not modelled	21.1	13	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> Aminoacid dehydrogenase-like, C-terminal domain
110	<a href="#">c2y0dB_</a>	Alignment	not modelled	21.0	17	<b>PDB header:</b> oxidoreductase <b>Chain:</b> B: <b>PDB Molecule:</b> udp-glucose dehydrogenase; <b>PDBTitle:</b> bcec mutation y10k
111	<a href="#">c2qytA_</a>	Alignment	not modelled	21.0	24	<b>PDB header:</b> oxidoreductase <b>Chain:</b> A: <b>PDB Molecule:</b> 2-dehydropantoate 2-reductase; <b>PDBTitle:</b> crystal structure of 2-dehydropantoate 2-reductase from porphyromonas2 gingivalis w83
112	<a href="#">d1jaya_</a>	Alignment	not modelled	20.4	37	<b>Fold:</b> NAD(P)-binding Rossmann-fold domains <b>Superfamily:</b> NAD(P)-binding Rossmann-fold domains <b>Family:</b> 6-phosphogluconate dehydrogenase-like, N-terminal domain
113	<a href="#">c1d4fD_</a>	Alignment	not modelled	20.2	21	<b>PDB header:</b> hydrolase <b>Chain:</b> D: <b>PDB Molecule:</b> s-adenosylhomocysteine hydrolase; <b>PDBTitle:</b> crystal structure of recombinant rat-liver d244e mutant s-2 adenosylhomocysteine hydrolase