













Phyre²

| | |
|---------------|--------------------------------|
| Email | i.a.kelley@imperial.ac.uk |
| Description | P31808 |
| Date | Thu Jan 5 11:48:45 GMT 2012 |
| Unique Job ID | bde535b825840ce9 |

Detailed template information

| # | Template | Alignment Coverage | 3D Model | Confidence | % i.d. | Template Information |
|----|-------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|------------|--------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1 | d1iy8a_ |  |  | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 2 | c3lf2B_ |  |  | 100.0 | 22 | PDB header: oxidoreductase Chain: B: PDB Molecule: short chain oxidoreductase q9hya2; PDBTitle: nadph bound structure of the short chain oxidoreductase q9hya2 from2 pseudomonas aeruginosa pao1 containing an atypical catalytic center |
| 3 | c3svtA_ |  |  | 100.0 | 26 | PDB header: oxidoreductase Chain: A: PDB Molecule: short-chain type dehydrogenase/reductase; PDBTitle: structure of a short-chain type dehydrogenase/reductase from2 mycobacterium ulcerans |
| 4 | d1w6ua_ |  |  | 100.0 | 18 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 5 | d1xhla_ |  |  | 100.0 | 24 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 6 | d1yxma1 |  |  | 100.0 | 15 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 7 | d2ae2a_ |  |  | 100.0 | 22 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 8 | d1aela_ |  |  | 100.0 | 25 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 9 | c3o38D_ |  |  | 100.0 | 21 | PDB header: oxidoreductase Chain: D: PDB Molecule: short chain dehydrogenase; PDBTitle: crystal structure of a short chain dehydrogenase from mycobacterium smegmatis |
| 10 | d2bgka1 |  |  | 100.0 | 22 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 11 | d1xkqa_ |  |  | 100.0 | 24 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |

| | | | | | | |
|----|-------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 12 | d1geea_ |  |  | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 13 | d1nffa_ |  |  | 100.0 | 24 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 14 | c2zatC_ |  |  | 100.0 | 26 | PDB header: oxidoreductase Chain: C: PDB Molecule: dehydrogenase/reductase sdr family member 4; PDBTitle: crystal structure of a mammalian reductase |
| 15 | c3rihB_ |  |  | 100.0 | 26 | PDB header: oxidoreductase Chain: B: PDB Molecule: short chain dehydrogenase or reductase; PDBTitle: crystal structure of a putative short chain dehydrogenase or reductase2 from mycobacterium abscessus |
| 16 | c3ctxA_ |  |  | 100.0 | 23 | PDB header: oxidoreductase Chain: A: PDB Molecule: dehydrogenase with different specificities; PDBTitle: quaternary complex structure of gluconate 5-dehydrogenase2 from streptococcus suis type 2 |
| 17 | c3nugA_ |  |  | 100.0 | 22 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-oxoacyl-(acyl-carrier protein) reductase; PDBTitle: crystal structure of wild type tetrameric pyridoxal 4-dehydrogenase2 from mesorhizobium loti |
| 18 | d1fmca_ |  |  | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 19 | d1zema1 |  |  | 100.0 | 24 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 20 | c2cfcb_ |  |  | 100.0 | 22 | PDB header: oxidoreductase Chain: B: PDB Molecule: 2-(r)-hydroxypropyl-com dehydrogenase; PDBTitle: structural basis for stereo selectivity in the (r)- and2 (s)-hydroxypropylethane thiosulfonate dehydrogenases |
| 21 | c3k31B_ |  | not modelled | 100.0 | 16 | PDB header: oxidoreductase Chain: B: PDB Molecule: enoyl-(acyl-carrier-protein) reductase; PDBTitle: crystal structure of enoyl-(acyl-carrier-protein) reductase from2 anaplasma phagocytophilum in complex with nad at 1.9a resolution |
| 22 | d1hdca_ |  | not modelled | 100.0 | 25 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 23 | c3toxG_ |  | not modelled | 100.0 | 24 | PDB header: oxidoreductase Chain: G: PDB Molecule: short chain dehydrogenase; PDBTitle: crystal structure of a short chain dehydrogenase in complex with2 nad(p) from sinorhizobium meliloti 1021 |
| 24 | c3r3sD_ |  | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: D: PDB Molecule: oxidoreductase; PDBTitle: structure of the ygha oxidoreductase from salmonella enterica |
| 25 | c3ijrF_ |  | not modelled | 100.0 | 19 | PDB header: oxidoreductase Chain: F: PDB Molecule: oxidoreductase, short chain dehydrogenase/reductase family; PDBTitle: 2.05 angstrom resolution crystal structure of a short chain2 dehydrogenase from bacillus anthracis str. 'ames ancestor' in complex3 with nad+ |
| 26 | c3gvcB_ |  | not modelled | 100.0 | 21 | PDB header: oxidoreductase Chain: B: PDB Molecule: probable short-chain type PDBTitle: crystal structure of probable short-chain dehydrogenase-2 reductase from mycobacterium tuberculosis |
| 27 | d1pr9a_ |  | not modelled | 100.0 | 22 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| | | | | | | Fold: NAD(P)-binding Rossmann-fold domains |

| | | | | | | |
|----|-------------------------|-----------|--------------|-------|----|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 28 | d1g0oa | Alignment | not modelled | 100.0 | 20 | Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 29 | c3t7cC | Alignment | not modelled | 100.0 | 29 | PDB header: oxidoreductase Chain: C: PDB Molecule: carveol dehydrogenase; PDBTitle: crystal structure of carveol dehydrogenase from mycobacterium avium2 bound to nad |
| 30 | c3imfA | Alignment | not modelled | 100.0 | 20 | PDB header: oxidoreductase Chain: A: PDB Molecule: short chain dehydrogenase; PDBTitle: 1.99 angstrom resolution crystal structure of a short chain2 dehydrogenase from bacillus anthracis str. 'ames ancestor' |
| 31 | c2jjyD | Alignment | not modelled | 100.0 | 12 | PDB header: oxidoreductase Chain: D: PDB Molecule: enoyl-[acyl-carrier-protein] reductase; PDBTitle: crystal structure of francisella tularensis enoyl reductase2 (ftfab1) with bound nad |
| 32 | d1zk4a1 | Alignment | not modelled | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 33 | c1w4zA | Alignment | not modelled | 100.0 | 22 | PDB header: antibiotic biosynthesis Chain: A: PDB Molecule: ketoacyl reductase; PDBTitle: structure of actinorhodin polyketide (actii) reductase |
| 34 | c3itdA | Alignment | not modelled | 100.0 | 16 | PDB header: oxidoreductase Chain: A: PDB Molecule: 17beta-hydroxysteroid dehydrogenase; PDBTitle: crystal structure of an inactive 17beta-hydroxysteroid dehydrogenase2 (y167f mutated form) from fungus cochlidiobolus lunatus |
| 35 | c3icca | Alignment | not modelled | 100.0 | 24 | PDB header: oxidoreductase Chain: A: PDB Molecule: putative 3-oxoacyl-(acyl carrier protein) reductase; PDBTitle: crystal structure of a putative 3-oxoacyl-(acyl carrier protein)2 reductase from bacillus anthracis at 1.87 a resolution |
| 36 | d2rhca1 | Alignment | not modelled | 100.0 | 22 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 37 | c3ai3A | Alignment | not modelled | 100.0 | 21 | PDB header: oxidoreductase Chain: A: PDB Molecule: nadph-sorbose reductase; PDBTitle: the crystal structure of l-sorbose reductase from gluconobacter2 frateurii complexed with nadph and l-sorbose |
| 38 | c3pk0B | Alignment | not modelled | 100.0 | 25 | PDB header: oxidoreductase Chain: B: PDB Molecule: short-chain dehydrogenase/reductase sdr; PDBTitle: crystal structure of short-chain dehydrogenase/reductase sdr from2 mycobacterium smegmatis |
| 39 | c2z1nA | Alignment | not modelled | 100.0 | 16 | PDB header: oxidoreductase Chain: A: PDB Molecule: dehydrogenase; PDBTitle: crystal structure of ape0912 from aeropyrum pernix k1 |
| 40 | d1k2wa | Alignment | not modelled | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 41 | c3ak4C | Alignment | not modelled | 100.0 | 26 | PDB header: oxidoreductase Chain: C: PDB Molecule: nadh-dependent quinuclidinone reductase; PDBTitle: crystal structure of nadh-dependent quinuclidinone reductase from2 agrobacterium tumefaciens |
| 42 | d2c07a1 | Alignment | not modelled | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 43 | c2c07A | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-oxoacyl-(acyl-carrier protein) reductase; PDBTitle: oxoacyl-acp reductase of plasmodium falciparum |
| 44 | c2pd6D | Alignment | not modelled | 100.0 | 24 | PDB header: oxidoreductase Chain: D: PDB Molecule: estradiol 17-beta-dehydrogenase 8; PDBTitle: structure of human hydroxysteroid dehydrogenase type 8, hsd17b8 |
| 45 | d1x1ta1 | Alignment | not modelled | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 46 | d1q7ba | Alignment | not modelled | 100.0 | 25 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 47 | c3uveC | Alignment | not modelled | 100.0 | 19 | PDB header: oxidoreductase Chain: C: PDB Molecule: carveol dehydrogenase ((+)-trans-carveol dehydrogenase); PDBTitle: crystal structure of carveol dehydrogenase ((+)-trans-carveol2 dehydrogenase) from mycobacterium avium |
| 48 | c3emkA | Alignment | not modelled | 100.0 | 27 | PDB header: oxidoreductase Chain: A: PDB Molecule: glucose/ribitol dehydrogenase; PDBTitle: 2.5a crystal structure of glucose/ribitol dehydrogenase2 from brucella melitensis |
| 49 | c3ek2D | Alignment | not modelled | 100.0 | 16 | PDB header: oxidoreductase Chain: D: PDB Molecule: enoyl-(acyl-carrier-protein) reductase (nadh); PDBTitle: crystal structure of enoyl-(acyl carrier protein) reductase2 from burkholderia pseudomallei 1719b |
| 50 | d1hxha | Alignment | not modelled | 100.0 | 20 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 51 | d2pd4a1 | Alignment | not modelled | 100.0 | 17 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 52 | c3rkuC | Alignment | not modelled | 100.0 | 25 | PDB header: oxidoreductase Chain: C: PDB Molecule: oxidoreductase ymr226c; PDBTitle: substrate fingerprint and the structure of nadp+-dependent serine2 dehydrogenase from saccharomyces cerevisiae complexed with nadp+ |

| | | | | | | |
|----|-------------------------|-----------|--------------|-------|----|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 53 | c3grkE | Alignment | not modelled | 100.0 | 16 | PDB header: oxidoreductase Chain: E: PDB Molecule: enoyl-(acyl-carrier-protein) reductase (nadh); PDBTitle: crystal structure of short chain dehydrogenase reductase2 sdr glucose-ribitol dehydrogenase from brucella melitensis |
| 54 | d1ydea1 | Alignment | not modelled | 100.0 | 26 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 55 | c2p91A | Alignment | not modelled | 100.0 | 15 | PDB header: oxidoreductase Chain: A: PDB Molecule: enoyl-[acyl-carrier-protein] reductase [nadh]; PDBTitle: crystal structure of enoyl-[acyl-carrier-protein] reductase (nadh)2 from aquifex aeolicus vf5 |
| 56 | d2ew8a1 | Alignment | not modelled | 100.0 | 20 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 57 | d1ulua | Alignment | not modelled | 100.0 | 18 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 58 | d1bdba | Alignment | not modelled | 100.0 | 17 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 59 | c3uf0A | Alignment | not modelled | 100.0 | 30 | PDB header: oxidoreductase Chain: A: PDB Molecule: short-chain dehydrogenase/reductase sdr; PDBTitle: crystal structure of a putative nad(p) dependent gluconate 5-2 dehydrogenase from beutenbergia cavernae(efi target efi-502044) with3 bound nadp (low occupancy) |
| 60 | c3ftpD | Alignment | not modelled | 100.0 | 26 | PDB header: oxidoreductase Chain: D: PDB Molecule: 3-oxoacyl-[acyl-carrier protein] reductase; PDBTitle: crystal structure of 3-ketoacyl-(acyl-carrier-protein)2 reductase from burkholderia pseudomallei at 2.05 a3 resolution |
| 61 | c3gr6A | Alignment | not modelled | 100.0 | 15 | PDB header: oxidoreductase Chain: A: PDB Molecule: enoyl-[acyl-carrier-protein] reductase [nadh]; PDBTitle: crystal structure of the staphylococcus aureus enoyl-acyl2 carrier protein reductase (fabi) in complex with nadp and triclosan |
| 62 | c3sjuA | Alignment | not modelled | 100.0 | 20 | PDB header: oxidoreductase Chain: A: PDB Molecule: keto reductase; PDBTitle: hedamycin polyketide ketoreductase bound to nadph |
| 63 | c3oecA | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: A: PDB Molecule: carveol dehydrogenase (mytha.01326.c, a0r518 homolog); PDBTitle: crystal structure of carveol dehydrogenase from mycobacterium2 thermoresistibile |
| 64 | d1ja9a | Alignment | not modelled | 100.0 | 20 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 65 | c3i1jB | Alignment | not modelled | 100.0 | 48 | PDB header: oxidoreductase Chain: B: PDB Molecule: oxidoreductase, short chain PDBTitle: structure of a putative short chain dehydrogenase from2 pseudomonas syringae |
| 66 | c3pgxB | Alignment | not modelled | 100.0 | 25 | PDB header: oxidoreductase Chain: B: PDB Molecule: carveol dehydrogenase; PDBTitle: crystal structure of a putative carveol dehydrogenase from2 mycobacterium paratuberculosis bound to nicotinamide adenine3 dinucleotide |
| 67 | d2d1ya1 | Alignment | not modelled | 100.0 | 24 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 68 | c3v2gA | Alignment | not modelled | 100.0 | 20 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-oxoacyl-[acyl-carrier-protein] reductase; PDBTitle: crystal structure of a dehydrogenase/reductase from sinorhizobium2 meliloti 1021 |
| 69 | c3r1iB | Alignment | not modelled | 100.0 | 21 | PDB header: oxidoreductase Chain: B: PDB Molecule: short-chain type dehydrogenase/reductase; PDBTitle: crystal structure of a short-chain type dehydrogenase/reductase from2 mycobacterium marinum |
| 70 | d1gega | Alignment | not modelled | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 71 | d1ulsa | Alignment | not modelled | 100.0 | 24 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 72 | c2q2qG | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: G: PDB Molecule: beta-d-hydroxybutyrate dehydrogenase; PDBTitle: structure of d-3-hydroxybutyrate dehydrogenase from pseudomonas putida |
| 73 | c2uvdE | Alignment | not modelled | 100.0 | 25 | PDB header: oxidoreductase Chain: E: PDB Molecule: 3-oxoacyl-(acyl-carrier-protein) reductase; PDBTitle: the crystal structure of a 3-oxoacyl-(acyl carrier protein)2 reductase from bacillus anthracis (ba3989) |
| 74 | d1h5qa | Alignment | not modelled | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 75 | c3gk3D | Alignment | not modelled | 100.0 | 21 | PDB header: oxidoreductase Chain: D: PDB Molecule: acetoacetyl-coa reductase; PDBTitle: crystal structure of acetoacetyl-coa reductase from2 burkholderia pseudomallei 1710b |
| | | | | | | PDB header: oxidoreductase |

| | | | | | | |
|----|-------------------------|-----------|--------------|-------|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 76 | c2p68A | Alignment | not modelled | 100.0 | 26 | Chain: A: PDB Molecule: 3-oxoacyl-[acyl-carrier-protein] reductase; PDBTitle: crystal structure of aq_1716 from aquifex aeolicus vf5 |
| 77 | d1qsga | Alignment | not modelled | 100.0 | 16 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 78 | c3bmra | Alignment | not modelled | 100.0 | 15 | PDB header: oxidoreductase Chain: A: PDB Molecule: pteridine reductase; PDBTitle: structure of pteridine reductase 1 (ptr1) from trypanosoma2 brucei in ternary complex with cofactor (nadp+) and3 inhibitor (compound ax6) |
| 79 | d1spxa | Alignment | not modelled | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 80 | d1o5ia | Alignment | not modelled | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 81 | d1xg5a | Alignment | not modelled | 100.0 | 18 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 82 | c3n74A | Alignment | not modelled | 100.0 | 19 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-ketoacyl-(acyl-carrier-protein) reductase; PDBTitle: crystal structure of 3-ketoacyl-(acyl-carrier-protein) reductasefrom2 brucella melitensis |
| 83 | c3f5sA | Alignment | not modelled | 100.0 | 99 | PDB header: oxidoreductase Chain: A: PDB Molecule: dehydrogenase; PDBTitle: crystal structure of putative short chain dehydrogenase from shigella2 flexneri 2a str. 301 |
| 84 | c2gioA | Alignment | not modelled | 100.0 | 17 | PDB header: unknown function Chain: A: PDB Molecule: enoyl-(acyl-carrier-protein) reductase; PDBTitle: x-ray structure of enoyl-acyl carrier protein reductase from bacillus2 anthracis with triclosan |
| 85 | c3lylB | Alignment | not modelled | 100.0 | 22 | PDB header: oxidoreductase Chain: B: PDB Molecule: 3-oxoacyl-(acyl-carrier-protein) reductase; PDBTitle: structure of 3-oxoacyl-acylcarrier protein reductase, fabg2 from francisella tularensis |
| 86 | c3afnC | Alignment | not modelled | 100.0 | 19 | PDB header: oxidoreductase Chain: C: PDB Molecule: carbonyl reductase; PDBTitle: crystal structure of aldose reductase a1-r complexed with nadp |
| 87 | d1vl8a | Alignment | not modelled | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 88 | c2b4qB | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: B: PDB Molecule: rharnolipids biosynthesis 3-oxoacyl-[acyl- PDBTitle: pseudomonas aeruginosa rhlg/nadp active-site complex |
| 89 | c3i4fD | Alignment | not modelled | 100.0 | 14 | PDB header: oxidoreductase Chain: D: PDB Molecule: 3-oxoacyl-[acyl-carrier protein] reductase; PDBTitle: structure of putative 3-oxoacyl-reductase from bacillus thuringiensis |
| 90 | c3gdfa | Alignment | not modelled | 100.0 | 17 | PDB header: oxidoreductase Chain: A: PDB Molecule: probable nadp-dependent mannitol dehydrogenase; PDBTitle: crystal structure of the nadp-dependent mannitol dehydrogenase from2 cladosporium herbarum. |
| 91 | d2a4ka1 | Alignment | not modelled | 100.0 | 23 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 92 | c3v2hb | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: B: PDB Molecule: d-beta-hydroxybutyrate dehydrogenase; PDBTitle: the crystal structure of d-beta-hydroxybutyrate dehydrogenase from2 sinorhizobium meliloti |
| 93 | c3ppiA | Alignment | not modelled | 100.0 | 17 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-hydroxyacyl-coa dehydrogenase type-2; PDBTitle: crystal structure of 3-hydroxyacyl-coa dehydrogenase type-2 from2 mycobacterium avium |
| 94 | c3uxyC | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: C: PDB Molecule: short-chain dehydrogenase/reductase sdr; PDBTitle: the crystal structure of short chain dehydrogenase from rhodobacter2 sphaeroides |
| 95 | c3sx2F | Alignment | not modelled | 100.0 | 22 | PDB header: oxidoreductase Chain: F: PDB Molecule: putative 3-ketoacyl-(acyl-carrier-protein) reductase; PDBTitle: crystal structure of a putative 3-ketoacyl-(acyl-carrier-protein)2 reductase from mycobacterium paratuberculosis in complex with nad |
| 96 | d1e6wa | Alignment | not modelled | 100.0 | 16 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 97 | c2ph3B | Alignment | not modelled | 100.0 | 23 | PDB header: oxidoreductase Chain: B: PDB Molecule: 3-oxoacyl-[acyl carrier protein] reductase; PDBTitle: crystal structure of 3-oxoacyl-[acyl carrier protein] reductase2 ttha0415 from thermus thermophilus |
| 98 | d1cyda | Alignment | not modelled | 100.0 | 21 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 99 | c3diIB | Alignment | not modelled | 100.0 | 21 | PDB header: oxidoreductase Chain: B: PDB Molecule: short-chain dehydrogenase/reductase sdr; PDBTitle: crystal structure of a carbohydrate specific scor |

| | | | | | |
|-----|-------------------------|-----------|--------------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| | | | | | enzyme2 from clostridium thermocellum, ligand-free form |
| 100 | c2qhxB | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: B: PDB Molecule: pteridine reductase 1; PDBTitle: structure of pteridine reductase from leishmania major2 complexed with a ligand |
| 101 | c3oidA | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: A: PDB Molecule: enoyl-[acyl-carrier-protein] reductase [nadph]; PDBTitle: crystal structure of enoyl-acp reductases iii (fabl) from b. subtilis2 (complex with nadp and tcl) |
| 102 | d1mxha | Alignment | not modelled | 100.0 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 103 | d1p33a | Alignment | not modelled | 100.0 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 104 | c3ezIA | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: A: PDB Molecule: acetoacetyl-coa reductase; PDBTitle: crystal structure of acetylacetyl-coa reductase from2 burkholderia pseudomallei 1710b |
| 105 | d1edoA | Alignment | not modelled | 100.0 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 106 | c2wdzD | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: D: PDB Molecule: short-chain dehydrogenase/reductase; PDBTitle: crystal structure of the short chain dehydrogenase2 galactitol-dehydrogenase (gatdh) of rhodobacter3 sphaerooides in complex with nad+ and 1,2-pentandiol |
| 107 | c3sc4A | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: A: PDB Molecule: short chain dehydrogenase (a0qtm2 homolog); PDBTitle: crystal structure of a short chain dehydrogenase (a0qtm2 homolog)2 mycobacterium thermoeristibile |
| 108 | c2dteB | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: B: PDB Molecule: glucose 1-dehydrogenase related protein; PDBTitle: structure of thermoplasma acidophilum aldehydoxide dehydrogenase (aldt2) in complex with nadh |
| 109 | c3t4xA | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: A: PDB Molecule: oxidoreductase, short chain dehydrogenase/reductase family; PDBTitle: short chain dehydrogenase/reductase family oxidoreductase from2 bacillus anthracis str. ames ancestor |
| 110 | d2h7ma1 | Alignment | not modelled | 100.0 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 111 | d2ag5a1 | Alignment | not modelled | 100.0 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 112 | d2gdza1 | Alignment | not modelled | 100.0 | Fold: NAD(P)-binding Rossmann-fold domains Superfamily: NAD(P)-binding Rossmann-fold domains Family: Tyrosine-dependent oxidoreductases |
| 113 | c3e03C | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: C: PDB Molecule: short chain dehydrogenase; PDBTitle: crystal structure of a putative dehydrogenase from2 xanthomonas campestris |
| 114 | c3ctmH | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: H: PDB Molecule: carbonyl reductase; PDBTitle: crystal structure of a carbonyl reductase from candida2 parapsilosis with anti-prelog stereo-specificity |
| 115 | c2nm0B | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: B: PDB Molecule: probable 3-oxacyl-(acyl-carrier-protein) reductase; PDBTitle: crystal structure of sco1815: a beta-ketoacyl-acyl carrier protein2 reductase from streptomyces coelicolor a3(2) |
| 116 | c3f9iB | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: B: PDB Molecule: 3-oxoacyl-[acyl-carrier-protein] reductase; PDBTitle: crystal structure of 3-ketoacyl-(acyl-carrier-protein) reductase2 rickettsia prowazekii |
| 117 | c3osuA | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-oxoacyl-[acyl-carrier-protein] reductase; PDBTitle: crystal structure of the 3-oxoacyl-acyl carrier protein reductase,2 fabg, from staphylococcus aureus |
| 118 | c3llsB | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: B: PDB Molecule: 3-ketoacyl-(acyl-carrier-protein) reductase; PDBTitle: crystal structure of 3-ketoacyl-(acyl-carrier-protein) reductase from2 mycobacterium tuberculosis |
| 119 | c3ksuA | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: A: PDB Molecule: 3-oxoacyl-acyl carrier protein reductase; PDBTitle: crystal structure of short-chain dehydrogenase from2 oenococcus oeni psu-1 |
| 120 | c2fwmX | Alignment | not modelled | 100.0 | PDB header: oxidoreductase Chain: X: PDB Molecule: 2,3-dihydro-2,3-dihydroxybenzoate dehydrogenase; PDBTitle: crystal structure of e. coli enta, a 2,3-dihydrodihydroxy benzoate2 dehydrogenase |