












# Phyre<sup>2</sup>

Email	i.a.kelley@imperial.ac.uk
Description	P0A9X9
Date	Thu Jan 5 11:11:39 GMT 2012
Unique Job ID	d61a380c7dd9785c

Detailed template information

#	Template	Alignment Coverage	3D Model	Confidence	% i.d.	Template Information
1	<a href="#">d1h95a_</a>			99.9	44	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
2	<a href="#">d1mica_</a>			99.9	100	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
3	<a href="#">d2es2a1</a>			99.9	63	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
4	<a href="#">d1c9oa_</a>			99.9	58	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
5	<a href="#">c3camB_</a>			99.9	61	<b>PDB header:</b> gene regulation <b>Chain:</b> B: <b>PDB Molecule:</b> cold-shock domain family protein; <b>PDBTitle:</b> crystal structure of the cold shock domain protein from neisseria2 meningitidis
6	<a href="#">c3a0jB_</a>			99.9	54	<b>PDB header:</b> transcription <b>Chain:</b> B: <b>PDB Molecule:</b> cold shock protein; <b>PDBTitle:</b> crystal structure of cold shock protein 1 from thermus2 thermophilus hb8
7	<a href="#">d1g6pa_</a>			99.9	57	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
8	<a href="#">c2k5nA_</a>			99.9	26	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> putative cold-shock protein; <b>PDBTitle:</b> solution nmr structure of the n-terminal domain of protein2 eca1580 from erwinia carotovora, northeast structural3 genomics consortium target ewr156a
9	<a href="#">c2kcmA_</a>			99.9	34	<b>PDB header:</b> nucleic acid binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> cold shock domain family protein; <b>PDBTitle:</b> solution nmr structure of the n-terminal ob-domain of so_1732 from2 shewanella oneidensis. northeast structural genomics consortium3 target sor210a.
10	<a href="#">c3aqqD_</a>			99.8	34	<b>PDB header:</b> dna binding protein <b>Chain:</b> D: <b>PDB Molecule:</b> calcium-regulated heat stable protein 1; <b>PDBTitle:</b> crystal structure of human crhsp-24
11	<a href="#">c3trzE_</a>			99.8	46	<b>PDB header:</b> rna binding protein/rna <b>Chain:</b> E: <b>PDB Molecule:</b> protein lin-28 homolog a; <b>PDBTitle:</b> mouse lin28a in complex with let-7d microRNA pre-element

12	<a href="#">c2ytyA_</a>			99.8	29	<b>PDB header:</b> rna binding protein <b>Chain:</b> A; <b>PDB Molecule:</b> cold shock domain-containing protein e1; <b>PDBTitle:</b> solution structure of the fourth cold-shock domain of the human2 kiaa0885 protein (unr protein)
13	<a href="#">c2ytxA_</a>			99.7	31	<b>PDB header:</b> rna binding protein <b>Chain:</b> A; <b>PDB Molecule:</b> cold shock domain-containing protein e1; <b>PDBTitle:</b> solution structure of the second cold-shock domain of the human2 kiaa0885 protein (unr protein)
14	<a href="#">d1wfqa_</a>			99.7	21	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
15	<a href="#">c1x65A_</a>			99.7	34	<b>PDB header:</b> rna binding protein <b>Chain:</b> A; <b>PDB Molecule:</b> unr protein; <b>PDBTitle:</b> solution structure of the third cold-shock domain of the human2 kiaa0885 protein (unr protein)
16	<a href="#">c2bh8B_</a>			99.6	70	<b>PDB header:</b> transcription <b>Chain:</b> B; <b>PDB Molecule:</b> 1b11; <b>PDBTitle:</b> combinatorial protein 1b11
17	<a href="#">c2ytvA_</a>			99.6	32	<b>PDB header:</b> rna binding protein <b>Chain:</b> A; <b>PDB Molecule:</b> cold shock domain-containing protein e1; <b>PDBTitle:</b> solution structure of the fifth cold-shock domain of the human2 kiaa0885 protein (unr protein)
18	<a href="#">d2ix0a2</a>			97.6	22	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
19	<a href="#">c2ix1A_</a>			97.2	25	<b>PDB header:</b> hydrolase <b>Chain:</b> A; <b>PDB Molecule:</b> exoribonuclease 2; <b>PDBTitle:</b> rnase ii d209n mutant
20	<a href="#">d1a62a2</a>			96.5	28	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
21	<a href="#">c2a8vA_</a>		not modelled	96.3	28	<b>PDB header:</b> protein/rna <b>Chain:</b> A; <b>PDB Molecule:</b> rna binding domain of rho transcription <b>PDBTitle:</b> rho transcription termination factor/rna complex
22	<a href="#">d1smxa_</a>		not modelled	95.7	20	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
23	<a href="#">d2ix0a1</a>		not modelled	93.9	18	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
24	<a href="#">c2k52A_</a>		not modelled	93.2	9	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A; <b>PDB Molecule:</b> uncharacterized protein mj1198; <b>PDBTitle:</b> structure of uncharacterized protein mj1198 from2 methanocaldococcus jannaschii. northeast structural3 genomics target mjr117b
25	<a href="#">d1kl9a2</a>		not modelled	92.8	11	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
26	<a href="#">c2vnuD_</a>		not modelled	91.9	19	<b>PDB header:</b> hydrolase/rna <b>Chain:</b> D; <b>PDB Molecule:</b> exosome complex exonuclease rrp44; <b>PDBTitle:</b> crystal structure of sc rrp44
27	<a href="#">d1y14b1</a>		not modelled	90.9	22	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
28	<a href="#">d1u0la1</a>		not modelled	90.6	12	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
						<b>PDB header:</b> gene regulation

29	<a href="#">c3go5A</a>		not modelled	89.4	16	<b>Chain:</b> A: <b>PDB Molecule:</b> multidomain protein with s1 rna-binding domains; <b>PDBTitle:</b> crystal structure of a multidomain protein with nucleic acid binding2 domains (sp_0946) from streptococcus pneumoniae tigr4 at 1.40 a3 resolution
30	<a href="#">d1hh2p1</a>		not modelled	87.5	15	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
31	<a href="#">c1xpuB</a>		not modelled	85.4	34	<b>PDB header:</b> transcription/rna <b>Chain:</b> B: <b>PDB Molecule:</b> rho transcription termination factor; <b>PDBTitle:</b> structural mechanism of inhibition of the rho transcription2 termination factor by the antibiotic 5a-(3-formylphenylsulfanyl)-3 dihydrobicyclomycin (fpdb)
32	<a href="#">d2c35b1</a>		not modelled	85.0	18	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
33	<a href="#">c2cqoA</a>		not modelled	84.9	23	<b>PDB header:</b> ribosome <b>Chain:</b> A: <b>PDB Molecule:</b> nucleolar protein of 40 kda; <b>PDBTitle:</b> solution structure of the s1 rna binding domain of human2 hypothetical protein flj11067
34	<a href="#">c1hh2P</a>		not modelled	84.6	15	<b>PDB header:</b> transcription regulation <b>Chain:</b> P: <b>PDB Molecule:</b> n utilization substance protein a; <b>PDBTitle:</b> crystal structure of nusa from thermotoga maritima
35	<a href="#">c1l2fA</a>		not modelled	81.7	15	<b>PDB header:</b> transcription <b>Chain:</b> A: <b>PDB Molecule:</b> n utilization substance protein a; <b>PDBTitle:</b> crystal structure of nusa from thermotoga maritima: a2 structure-based role of the n-terminal domain
36	<a href="#">d2z0sa1</a>		not modelled	81.6	10	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
37	<a href="#">c2oceA</a>		not modelled	81.2	19	<b>PDB header:</b> structural genomics, unknown function <b>Chain:</b> A: <b>PDB Molecule:</b> hypothetical protein pa5201; <b>PDBTitle:</b> crystal structure of tex family protein pa5201 from2 pseudomonas aeruginosa
38	<a href="#">d1wi5a</a>		not modelled	80.3	17	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
39	<a href="#">c2khiA</a>		not modelled	77.6	22	<b>PDB header:</b> ribosomal protein <b>Chain:</b> A: <b>PDB Molecule:</b> 30s ribosomal protein s1; <b>PDBTitle:</b> nmr structure of the domain 4 of the e. coli ribosomal2 protein s1
40	<a href="#">c1u0IB</a>		not modelled	77.5	14	<b>PDB header:</b> hydrolase <b>Chain:</b> B: <b>PDB Molecule:</b> probable gtpase engc; <b>PDBTitle:</b> crystal structure of yeq from thermotoga maritima
41	<a href="#">d2nn6h1</a>		not modelled	77.2	13	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
42	<a href="#">c2I55A</a>		not modelled	76.6	21	<b>PDB header:</b> metal binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> silb,silver efflux protein, mfp component of the three <b>PDBTitle:</b> solution structure of the c-terminal domain of silb from cupriavidus2 metallidurans
43	<a href="#">d3bzka4</a>		not modelled	74.0	19	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
44	<a href="#">c2wp8J</a>		not modelled	73.8	17	<b>PDB header:</b> hydrolase <b>Chain:</b> J: <b>PDB Molecule:</b> exosome complex exonuclease dis3; <b>PDBTitle:</b> yeast rrp44 nuclease
45	<a href="#">c2khjA</a>		not modelled	72.7	16	<b>PDB header:</b> ribosomal protein <b>Chain:</b> A: <b>PDB Molecule:</b> 30s ribosomal protein s1; <b>PDBTitle:</b> nmr structure of the domain 6 of the e. coli ribosomal2 protein s1
46	<a href="#">d2ba0a1</a>		not modelled	70.8	17	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
47	<a href="#">c2c4rL</a>		not modelled	70.8	18	<b>PDB header:</b> hydrolase <b>Chain:</b> L: <b>PDB Molecule:</b> ribonuclease e; <b>PDBTitle:</b> catalytic domain of e. coli rnase e
48	<a href="#">c1zeqX</a>		not modelled	69.0	17	<b>PDB header:</b> metal binding protein <b>Chain:</b> X: <b>PDB Molecule:</b> cation efflux system protein cufc; <b>PDBTitle:</b> 1.5 a structure of apo-cufc residues 6-88 from escherichia2 coli
49	<a href="#">d1q46a2</a>		not modelled	68.8	9	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
50	<a href="#">c1q46A</a>		not modelled	67.2	9	<b>PDB header:</b> translation <b>Chain:</b> A: <b>PDB Molecule:</b> translation initiation factor 2 alpha subunit; <b>PDBTitle:</b> crystal structure of the eif2 alpha subunit from2 saccharomyces cerevisiae
51	<a href="#">d1go3el</a>		not modelled	62.8	25	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
52	<a href="#">c2ahob</a>		not modelled	62.2	22	<b>PDB header:</b> translation <b>Chain:</b> B: <b>PDB Molecule:</b> translation initiation factor 2 alpha subunit; <b>PDBTitle:</b> structure of the archaeal initiation factor eif2 alpha-2 gamma heterodimer from sulfolobus solfataricus complexed3 with gdnpn
53	<a href="#">c3I0oB</a>		not modelled	61.0	31	<b>PDB header:</b> hydrolase <b>Chain:</b> B: <b>PDB Molecule:</b> transcription termination factor rho; <b>PDBTitle:</b> structure of rna-free rho transcription termination factor from2 thermotoga maritima
						<b>PDB header:</b> rna binding protein

54	<a href="#">c2k4KA</a>	Alignment	not modelled	60.3	19	<b>Chain:</b> A: <b>PDB Molecule:</b> general stress protein 13; <b>PDBTitle:</b> solution structure of gsp13 from bacillus subtilis
55	<a href="#">c1q8kA</a>	Alignment	not modelled	57.0	11	<b>PDB header:</b> translation <b>Chain:</b> A: <b>PDB Molecule:</b> eukaryotic translation initiation factor 2 <b>PDBTitle:</b> solution structure of alpha subunit of human eif2
56	<a href="#">c3h0gS</a>	Alignment	not modelled	57.0	28	<b>PDB header:</b> transcription <b>Chain:</b> S: <b>PDB Molecule:</b> dna-directed rna polymerase ii subunit rpb7; <b>PDBTitle:</b> rna polymerase ii from schizosaccharomyces pombe
57	<a href="#">d2je6i1</a>	Alignment	not modelled	56.2	19	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
58	<a href="#">c3psiA</a>	Alignment	not modelled	55.2	11	<b>PDB header:</b> transcription <b>Chain:</b> A: <b>PDB Molecule:</b> transcription elongation factor spt6; <b>PDBTitle:</b> crystal structure of the spt6 core domain from saccharomyces2 cerevisiae, form spt6(239-1451)
59	<a href="#">c2z0sA</a>	Alignment	not modelled	50.1	10	<b>PDB header:</b> rna binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> probable exosome complex rna-binding protein 1; <b>PDBTitle:</b> crystal structure of putative exosome complex rna-binding2 protein
60	<a href="#">c2eqsA</a>	Alignment	not modelled	48.6	19	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> atp-dependent rna helicase dhx8; <b>PDBTitle:</b> solution structure of the s1 rna binding domain of human2 atp-dependent rna helicase dhx8
61	<a href="#">d1h9ma2</a>	Alignment	not modelled	45.9	19	<b>Fold:</b> OB-fold <b>Superfamily:</b> MOP-like <b>Family:</b> BiMOP, duplicated molybdate-binding domain
62	<a href="#">d2ahob2</a>	Alignment	not modelled	45.6	22	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
63	<a href="#">c2c35F</a>	Alignment	not modelled	44.2	19	<b>PDB header:</b> polymerase <b>Chain:</b> F: <b>PDB Molecule:</b> dna-directed rna polymerase ii 19 kda <b>PDBTitle:</b> subunits rpb4 and rpb7 of human rna polymerase ii
64	<a href="#">c2xngA</a>	Alignment	not modelled	38.1	38	<b>PDB header:</b> rna binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> nuclear polyadenylated rna-binding protein 3; <b>PDBTitle:</b> structural insights into cis element recognition of non-2 polyadenylated rnas by the nab3-rrm
65	<a href="#">d1sroa</a>	Alignment	not modelled	34.6	31	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
66	<a href="#">c2i38A</a>	Alignment	not modelled	34.4	25	<b>PDB header:</b> rna binding protein/chimera <b>Chain:</b> A: <b>PDB Molecule:</b> fusion protein consists of immunoglobin g- <b>PDBTitle:</b> solution structure of the rrm of srp20
67	<a href="#">d1h9ma1</a>	Alignment	not modelled	32.5	11	<b>Fold:</b> OB-fold <b>Superfamily:</b> MOP-like <b>Family:</b> BiMOP, duplicated molybdate-binding domain
68	<a href="#">d1t9ha1</a>	Alignment	not modelled	31.6	20	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
69	<a href="#">c1yz6A</a>	Alignment	not modelled	31.1	22	<b>PDB header:</b> translation <b>Chain:</b> A: <b>PDB Molecule:</b> probable translation initiation factor 2 alpha <b>PDBTitle:</b> crystal structure of intact alpha subunit of aif2 from2 pyrococcus abyssi
70	<a href="#">c2qt7B</a>	Alignment	not modelled	30.7	28	<b>PDB header:</b> hydrolase <b>Chain:</b> B: <b>PDB Molecule:</b> receptor-type tyrosine-protein phosphatase-like <b>PDBTitle:</b> crystallographic structure of the mature ectodomain of the2 human receptor-type protein-tyrosine phosphatase ia-2 at3 1.30 angstroms
71	<a href="#">c2b8kG</a>	Alignment	not modelled	30.5	24	<b>PDB header:</b> transferase <b>Chain:</b> G: <b>PDB Molecule:</b> dna-directed rna polymerase ii 19 kda <b>PDBTitle:</b> 12-subunit rna polymerase ii
72	<a href="#">c1nt9G</a>	Alignment	not modelled	30.3	20	<b>PDB header:</b> transcription, transferase <b>Chain:</b> G: <b>PDB Molecule:</b> dna-directed rna polymerase ii 19 kd polypeptide; <b>PDBTitle:</b> complete 12-subunit rna polymerase ii
73	<a href="#">d1fr3a</a>	Alignment	not modelled	29.7	17	<b>Fold:</b> OB-fold <b>Superfamily:</b> MOP-like <b>Family:</b> Molybdate/tungstate binding protein MOP
74	<a href="#">c1j6qA</a>	Alignment	not modelled	29.7	22	<b>PDB header:</b> chaperone <b>Chain:</b> A: <b>PDB Molecule:</b> cytochrome c maturation protein e; <b>PDBTitle:</b> solution structure and characterization of the heme2 chaperone ccme
75	<a href="#">d1j6qa</a>	Alignment	not modelled	29.7	22	<b>Fold:</b> OB-fold <b>Superfamily:</b> Heme chaperone CcmE <b>Family:</b> Heme chaperone CcmE
76	<a href="#">c2dhxA</a>	Alignment	not modelled	28.1	21	<b>PDB header:</b> rna binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> poly (adp-ribose) polymerase family, member 10 <b>PDBTitle:</b> solution structure of the rrm domain in the human poly (adp-2 ribose) polymerase family, member 10 variant
77	<a href="#">d1guta</a>	Alignment	not modelled	27.0	11	<b>Fold:</b> OB-fold <b>Superfamily:</b> MOP-like <b>Family:</b> Molybdate/tungstate binding protein MOP
78	<a href="#">c2pmzE</a>	Alignment	not modelled	26.3	29	<b>PDB header:</b> translation, transferase <b>Chain:</b> E: <b>PDB Molecule:</b> dna-directed rna polymerase subunit e; <b>PDBTitle:</b> archaeal rna polymerase from sulfolobus solfataricus
79	<a href="#">d2vnud3</a>	Alignment	not modelled	26.1	17	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins

					<b>Family:</b> Cold shock DNA-binding domain-like
80	<a href="#">d2asba1</a>	Alignment	not modelled	24.7	<b>Fold:</b> OB-fold <b>Superfamily:</b> Nucleic acid-binding proteins <b>Family:</b> Cold shock DNA-binding domain-like
81	<a href="#">d1ueqa_</a>	Alignment	not modelled	23.6	<b>Fold:</b> PDZ domain-like <b>Superfamily:</b> PDZ domain-like <b>Family:</b> PDZ domain
82	<a href="#">d1tp5a1</a>	Alignment	not modelled	23.6	<b>Fold:</b> PDZ domain-like <b>Superfamily:</b> PDZ domain-like <b>Family:</b> PDZ domain
83	<a href="#">c1h9mB_</a>	Alignment	not modelled	22.7	<b>PDB header:</b> binding protein <b>Chain:</b> B: <b>PDB Molecule:</b> molybdenum-binding-protein; <b>PDBTitle:</b> two crystal structures of the cytoplasmic molybdate-binding2 protein modg suggest a novel cooperative binding mechanism3 and provide insights into ligand-binding specificity.4 peg-grown form with molybdate bound
84	<a href="#">c1go3E_</a>	Alignment	not modelled	21.7	<b>PDB header:</b> transferase <b>Chain:</b> E: <b>PDB Molecule:</b> dna-directed rna polymerase subunit e; <b>PDBTitle:</b> structure of an archaeal homolog of the eukaryotic rna2 polymerase ii rpb4/rpb7 complex
85	<a href="#">c2q3gA_</a>	Alignment	not modelled	21.1	<b>PDB header:</b> structural genomics <b>Chain:</b> A: <b>PDB Molecule:</b> pdz and lim domain protein 7; <b>PDBTitle:</b> structure of the pdz domain of human pdlim7 bound to a c-2 terminal extension from human beta-tropomyosin
86	<a href="#">c2je6l_</a>	Alignment	not modelled	20.5	<b>PDB header:</b> hydrolase <b>Chain:</b> I: <b>PDB Molecule:</b> exosome complex rna-binding protein 1; <b>PDBTitle:</b> structure of a 9-subunit archaeal exosome
87	<a href="#">c2asbA_</a>	Alignment	not modelled	20.0	<b>PDB header:</b> transcription/rna <b>Chain:</b> A: <b>PDB Molecule:</b> transcription elongation protein nusa; <b>PDBTitle:</b> structure of a mycobacterium tuberculosis nusa-rna complex
88	<a href="#">c2edzA_</a>	Alignment	not modelled	20.0	<b>PDB header:</b> signaling protein <b>Chain:</b> A: <b>PDB Molecule:</b> pdz domain-containing protein 1; <b>PDBTitle:</b> solution structures of the pdz domain of mus musculus pdz2 domain-containing protein 1
89	<a href="#">c2gu1A_</a>	Alignment	not modelled	19.8	<b>PDB header:</b> hydrolase <b>Chain:</b> A: <b>PDB Molecule:</b> zinc peptidase; <b>PDBTitle:</b> crystal structure of a zinc containing peptidase from vibrio cholerae
90	<a href="#">c2dc2A_</a>	Alignment	not modelled	19.7	<b>PDB header:</b> structural protein <b>Chain:</b> A: <b>PDB Molecule:</b> golgi associated pdz and coiled-coil motif <b>PDBTitle:</b> solution structure of pdz domain
91	<a href="#">c2opgB_</a>	Alignment	not modelled	19.7	<b>PDB header:</b> structural protein <b>Chain:</b> B: <b>PDB Molecule:</b> multiple pdz domain protein; <b>PDBTitle:</b> the crystal structure of the 10th pdz domain of mpdz
92	<a href="#">c3soeA_</a>	Alignment	not modelled	17.4	<b>PDB header:</b> signaling protein <b>Chain:</b> A: <b>PDB Molecule:</b> membrane-associated guanylate kinase, ww and pdz domain- <b>PDBTitle:</b> crystal structure of the 3rd pdz domain of the human membrane-2 associated guanylate kinase, ww and pdz domain-containing protein 33 (magi3)
93	<a href="#">d1qaua_</a>	Alignment	not modelled	17.3	<b>Fold:</b> PDZ domain-like <b>Superfamily:</b> PDZ domain-like <b>Family:</b> PDZ domain
94	<a href="#">c3ggeA_</a>	Alignment	not modelled	17.0	<b>PDB header:</b> protein binding <b>Chain:</b> A: <b>PDB Molecule:</b> pdz domain-containing protein gipc2; <b>PDBTitle:</b> crystal structure of the pdz domain of pdz domain-containing protein2 gipc2
95	<a href="#">c2iwnA_</a>	Alignment	not modelled	16.7	<b>PDB header:</b> signaling protein <b>Chain:</b> A: <b>PDB Molecule:</b> multiple pdz domain protein; <b>PDBTitle:</b> 3rd pdz domain of multiple pdz domain protein mpdz (casp2 target)
96	<a href="#">c2kvia_</a>	Alignment	not modelled	16.0	<b>PDB header:</b> rrna binding protein <b>Chain:</b> A: <b>PDB Molecule:</b> nuclear polyadenylated rna-binding protein 3; <b>PDBTitle:</b> structure of nab3 rrnm
97	<a href="#">c2i04B_</a>	Alignment	not modelled	15.5	<b>PDB header:</b> peptide binding protein <b>Chain:</b> B: <b>PDB Molecule:</b> membrane-associated guanylate kinase, ww and pdz <b>PDBTitle:</b> x-ray crystal structure of magi-1 pdz1 bound to the c-2 terminal peptide of hpv18 e6
98	<a href="#">c2kjda_</a>	Alignment	not modelled	15.5	<b>PDB header:</b> signaling protein <b>Chain:</b> A: <b>PDB Molecule:</b> sodium/hydrogen exchange regulatory cofactor nhe- <b>PDBTitle:</b> solution structure of extended pdz2 domain from nherf1 (150-2 270)
99	<a href="#">c3bpua_</a>	Alignment	not modelled	15.4	<b>PDB header:</b> transferase <b>Chain:</b> A: <b>PDB Molecule:</b> membrane-associated guanylate kinase, ww and pdz domain- <b>PDBTitle:</b> crystal structure of the 3rd pdz domain of human membrane associated2 guanylate kinase, c677s and c709s double mutant